

Xaletto – The economic management system for closed houses with straw bedding



Nutritional Value of Soy Products Fed to Pigs

DISEASE AND WELFARE HUSBANDRY FEED AND NUTRITION

by Jackie Linden
1 August 2013, at 12:00a.m.

A review of the nutrient composition of a range of soybean ingredients for pigs by H.H. Stein, J.A. Roth, K.M. Sotak and O.J. Rojas (University of Illinois, Urbana-Champaign) in the latest issue of 'Swine Focus'.

Soybeans is an important crop in the United States and are primarily used for animal feed, human food, and production of biofuels. Soybean meal (SBM) and other soy products contribute high-quality protein to diets fed to pigs because soy protein is rich in the limiting amino acids lysine, threonine and tryptophan that are present in relatively low concentrations in the most commonly fed cereal grains. Amino acids in soy protein also have a greater digestibility by pigs than amino acids in most other commonly fed protein sources.

Soy products are also a significant source of energy in diets fed to pigs and soybean meal contains as much digestible and metabolisable energy as corn. Although soy is usually fed to pigs in the form of soybean meal, full-fat soybeans may be included in the diets to increase the energy density of the diet. Due to the high fibre content of soy hulls, the energy concentration in de-hulled soybean products is greater than in products containing hulls.

The majority of the phosphorus in soy products is bound to phytic acid, which has a low digestibility to pigs but the digestibility of phosphorus in soy products may be increased to more than 60 per cent if diets are supplemented with microbial phytase.

Soybeans contain anti-nutritional factors that may limit growth of pigs but different technologies have been developed to reduce or eliminate the effects of anti-nutritional factors. All soybean products must be heat-treated prior to inclusion in swine diets to destroy trypsin inhibitors and lectins. Fermentation and enzyme treatment eliminate or greatly reduce allergenic proteins and

OUR SPONSORS



The Reference in Prevention for Animal Health



PARTNERS

[CID LINES](#), [Thermo Fisher Scientific](#), [Evonik](#), [Vencomatic Group](#), [Hypor](#), [Genesus](#), [PIC](#), [Novus International](#), [Adisseo](#)

Related news & articles >



13 May 2019

New survey highlights priority animal welfare issues in UK



7 May 2019

Trade disputes and weak financial performance wear on US farmers



3 May 2019

How can your vet help you to achieve sustainable production?



29 Apr 2019

Pig data meets big data: the precision pig farming revolution

oligosaccharides.

Oligosaccharides can also be reduced or eliminated by extracting the carbohydrates from soybean meal in an aqueous solution, which results in production of soy protein concentrate. Because of the low concentrations of oligosaccharides in fermented soybean meal, enzyme treated soybean meal, and soy protein concentrate, all of these products may be used in diets fed to weanling pigs.

Introduction

The United States is the world's largest producer of soybeans (*Glycine max*), and approximately 33 per cent of the world's soybeans are produced in the United States (ASA, 2012). In 2011, 75 million acres of soybeans were planted in the United States (Figure 1) and a total of 3,056 million bushels of soybeans were produced (ASA, 2012). Brazil produced 29 per cent of the world's soybeans followed by Argentina with 19 per cent (ASA, 2012). China, India, and Paraguay are other major producers of soybeans.




Figure 1. Top soybean-producing nations in 2011
(Values obtained from ASA, 2012)

Within the United States, Iowa is the largest producer of soybeans (Figure 2). In 2011, over 9.3 million acres of soybeans were planted in Iowa, and 466.12 million bushels of soybeans were produced. Illinois was the next largest producer with 416.42 million bushels, followed by Minnesota, Nebraska, Indiana, and Ohio (ASA, 2012).




Figure 2. Top soybean-producing states in 2011
(Values obtained from ASA, 2012).

The majority of soybeans are crushed into oil and soybean meal, and approximately 75 per cent of the soybean meal that is produced in the world is fed to pigs or poultry. In 2011, over 27 million short tons of soybean meal were fed to

livestock in the United States, with almost seven million short tons of soybean meal fed to pigs (ASA, 2012).

Soy Products Used in Swine Nutrition

Most soy products are fed to pigs in the form of soybean meal and its derivatives. These ingredients contain little fat because they are co-products of oil production, and thus, the fat has been removed.

Dehulled soybean meal contains less than two per cent fat on an as-fed basis, compared with whole full-fat soybeans that contain approximately 20 per cent fat (Table 1). Whole full-fat soybeans can be fed to increase the energy concentration of the diet, but because of the relatively high cost of soybean oil, this is usually not economical.

Table 1. Nutrient composition (per cent) of soybeans, soybean meal (SBM), and other soybean products (as-fed basis)¹

Product	Full-fat soybeans	Dehulled SBM	Non-dehulled SBM	Extruded-expelled SBM	Enzyme-treated SBM	Fermented SBM	Soy protein concentrate	Soy protein isolate
Dry matter	92.36	89.98	88.79	93.85	92.70	92.88	92.64	93.71
Crude protein	37.56	47.73	43.90	44.56	55.62	54.07	65.20	84.78
Ether extract	20.18	1.52	1.24	5.69	1.82	2.30	1.05	2.76
Carbohydrates and lignin	29.73	34.46	37.27	37.90	28.21	29.53	20.28	2.00
Ash	4.89	6.27	6.38	5.70	7.05	6.98	6.11	4.17

¹Values obtained from NRC (2012).

Swine producers commonly feed toasted soybean meal without further processing. However, soybean meal may be processed in different ways to increase the protein concentration and to deactivate anti-nutritional factors. Soybean meal is produced from whole or dehulled soybeans.

Dehulled soybean meal contains approximately 47.5 per cent crude protein on an as-fed basis, and is referred to as high-protein soybean meal. Non-dehulled soybean meal contains approximately 44 per cent crude protein and is referred to as low-protein soybean meal.

Conventional toasted soybean meal, or solvent-extracted dehulled soybean meal, is produced by extracting the fat from soy flour with a solvent, usually hexane, with a subsequent toasting step to remove residual hexane and to deactivate trypsin inhibitors and lectins.

* Fermentation or enzyme treatment of soybean meal eliminates the oligosaccharides in the meal

Expelled soybean meal is produced by extruding intact or dehulled soybeans followed by mechanical expelling of the oil. The extrusion-expelling process produces soybean meal with greater fat content than conventional soybean meal because the expelling process is less efficient than the

solvent extraction process in removing oil from the soybeans (Wang and Johnson, 2001). Therefore, extruded-expelled soybean meal contains between four and eight per cent fat, whereas solvent-extracted soybean meal contains less than two per cent fat.

Enzyme-treated soybean meal, a new product on the United States feed market, is produced by treating dehulled, solvent-extracted soybean meal for several hours with a proprietary blend of enzymes.

Enzyme treatment reduces the concentrations of oligosaccharides and allergenic proteins to create a product that can be fed to weanling pigs. To improve digestibility of phosphorus in enzyme-treated soybean meal, it may be treated with phytase (Goebel and Stein, 2011).

* Full fat soybeans contain 36 to 38 per cent crude protein and 19 to 20 per cent fat but dehulled soybean meal contains 46 to 48 per cent crude protein and less than two per cent fat.

Fermented soybean meal is produced by inoculating conventional soybean meal with the bacterium *Aspergillus oryzae* or other microbes. Fermenting the soybean meal eliminates the oligosaccharides and reduces the concentration of antigens in the meal (Cervantes-Pahm and Stein, 2010). Therefore, fermented soybean meal can be used instead of animal proteins in weanling pig diets without adversely affecting growth (Jones et al., 2010; Kim et al., 2010).

Soy protein concentrate is produced by aqueous ethanol extraction of water-soluble carbohydrates from defatted soybean meal. The ethanol extraction process removes the soluble carbohydrates, but the majority of the fibre in soybean meal is insoluble fibre and remains in soy protein concentrate. The resultant product contains at least 65 per cent crude protein (NRC, 2012).

Soy protein isolate is produced by solubilising the protein in soybean meal with water and precipitating the protein from the solution. This process removes the fat and carbohydrate components from the product (Cromwell, 2000); therefore, soy protein isolate contains at least 80 per cent crude protein (Middelbos and Fahey, 2008). The allergenic proteins, glycinin and β -conglycinin, are deactivated when soy protein concentrate and soy protein isolate are produced by extraction at temperatures greater than 50°C (Sissons et al, 1982) and both products are well tolerated by weanling pigs (Li et al., 1991). However, because of the high costs involved in producing soy protein isolate, this product is usually not used in the feeding of pigs.

Nutritional Value of Soybean Products

Energy

The energy concentration of soybeans and their derivatives depends on the concentration of fat, carbohydrates, and protein in the product. The digestible energy and

protein in the product. The digestible energy and metabolisable energy concentrations in full-fat soybeans are greater than in dehulled soybean meal (4,193 and 3,938 vs. 3,619 and 3,294 kcal/kg; Table 2). The reduced concentration of digestible and metabolisable energy in soybean meal compared with fullfat soybeans is a result of the reduced concentration of fat in soybean meal.

Table 2. Concentration of energy (kcal/kg) in soybeans, soybean meal (SBM), and other soybean products (as-fed basis)¹

Product	Full-fat soybeans	Dehulled SBM	Nondehulled SBM	Extruded-expelled SBM	Enzyme-treated SBM	Fermented SBM	Soy protein concentrate	Soy protein isolate
Gross energy	5227	4256	4257	4692	4451	4533	4605	5386
Digestible energy	4193	3619	3681	3876	3914	3975	4260	4150
Metabolizable energy	3938	3294	3382	3573	3536	3607	3817	3573
Net energy	2874	2087	2148	2344	-	-	2376	2187

¹Values obtained from NRC (2012).

Concentrations of digestible and metabolisable energy in enzyme-treated soybean meal are 3,914kcal per kg and 3,536kcal per kg, respectively. The energy content of enzyme-treated soybean meal is greater than that of conventional soybean meal because it has been further processed to remove antinutritional factors and antigenic proteins that decrease energy availability. The concentrations of digestible (3,975kcal per kg) and metabolisable (3,607kcal per kg) energy are also greater for fermented soybean meal than for conventional soybean meal because of the removal of oligosaccharides and other anti-nutritional factors.

Soy protein concentrate and soy protein isolate also have greater concentrations of energy than soybean meal because they contain a greater percentage of crude protein than soybean meal.

*

Dehulled soybean meal contains the same amount of digestible energy as corn.

Digestible energy values for soy protein concentrate and soy protein

isolate are 4,260kcal per kg and 4,150kcal per kg, respectively, and metabolisable energy values are 3,817 kcal per kg and 3,573 kcal per kg (NRC, 2012).

Carbohydrates

Soybeans are thought of primarily as a protein source but they also contain a total of 30 to 35 per cent carbohydrates, making soybeans a major carbohydrate contributor to the diet.

Soybeans contain two types of carbohydrates: structural and non-structural carbohydrates (Grieshop et al., 2003). The structural carbohydrates include cellulose and hemi-

cellulose, and the non-structural carbohydrates include sugars and oligosaccharides (Table 3).

Table 3. Concentrations (per cent) of carbohydrates and lignin in soybeans, soybean meal (SBM) and other soybean products (as-fed basis)¹


Product	Full-fat soybeans	Dehulled SBM	Nondehulled SBM	Extruded-expelled SBM	Soy protein concentrate	Soy protein isolate
Sucrose	6.42	4.30	7.63	7.1	0.67	0.13
Raffinose	0.77	3.78	0.90	0.77	0.46	-
Stachyose	3.89	7.33	4.32	4.88	0.91	-
Starch	1.89	1.89	1.89	1.89	1.89	1.89
Lignin	-	1.10	-	-	-	-
ADF	6.17	5.28	6.66	7.35	-	0.00
NDF	10.00	8.21	9.82	13.84	8.10	0.19

¹Values obtained from NRC (2012).

Oligosaccharides are considered anti-nutritional factors for young pigs because they reduce pig growth performance and increase the incidence of diarrhoea in weanling pigs.

Therefore, weanling pigs do not tolerate conventional soybean meal very well. However, most oligosaccharides are removed from fermented soybean meal and enzyme-treated soybean meal (Cervantes-Pahm and Stein, 2010) and these ingredients may, therefore, be included in diets fed to weanling pigs (Jones et al., 2010; Kim et al., 2010). Soy protein concentrate and soy protein isolate also have reduced concentrations of oligosaccharides compared with soybean meal because the water-soluble carbohydrates have been removed and these products may, therefore, also be included in diets fed to weanling pigs. Low-oligosaccharide soybean varieties are also being developed but are not commercially available at this time.

Dietary fibre has a reduced energy value compared with the energy of other nutrients. fibre also may negatively influence the digestibility and absorption of other nutrients and will contribute to a faster rate of passage in the intestinal tract of pigs. Pigs do not ferment dietary fibre very well; therefore, ingredients with a high fibre concentration have reduced dry matter digestibility compared with ingredients containing less fibre. However, among plant protein ingredients, dehulled soybean meal has the least dietary fibre content determined as the concentration of crude fibre, acid detergent fibre (ADF) or neutral detergent fibre (NDF; Figure 3). A consequence of the low concentration of fibre in soybean meal is that the energy concentration in soybean meal is similar to that of corn.






Figure 3. Fibre content of feed ingredients, per cent
(Values obtained from NRC, 2012)

Minerals

Most of the phosphorus in soybeans is bound to phytic acid (Table 4). Pigs cannot utilise most of the phytic acid bound phosphorus because they produce little intestinal phytase.

Table 4. Calcium and phosphorus concentrations and apparent (ATTD) and standardised (STTD) total tract digestibility of phosphorus in soybeans, soybean meal (SBM), and other soybean products (as-fed basis)¹

Product	Full-fat soybeans	Dehulled SBM	Nondehulled SBM	Enzyme treated SBM	Fermented SBM	Soy protein concentrate	Soy protein isolate
Total P, %	0.53	.71	.064	0.75	0.80	0.82	0.75
ATTD P, %	39	39	39	60	60	39	39
STTD P, %	48	48	48	66	66	48	48
Phytate bound P, %	0.33	0.38	0.36	-	-	-	-
Phytate bound P, % of total P	62.3	53.5	56.3	-	-	-	-
Non-phytate P, %	0.2	0.33	0.28	-	-	-	-
Non-phytate bound P, % of total P	37.7	46.5	43.8	-	-	-	-
Total Ca, %	0.31	0.33	0.35	0.31	0.3	0.32	0.17
ATTD Ca, %	-	62.9	-	60.9	50.7	-	-

¹Values obtained from NRC (2012).

Therefore, to make sure that there is enough digestible phosphorus in the diet, it is often necessary to add supplemental phosphorus in the form of monocalcium phosphate or dicalcium phosphate, which increases diet costs. Because the pig cannot utilise the phytic acid bound phosphorus, it is excreted in the pig's manure. Phosphorus run-off from the manure into aquatic ecosystems can lead to algal blooms and fish kills. Phytic acid also binds calcium, decreasing the amount of available calcium in the diet.

Supplementation of the diet with microbial phytase improves

phosphorus digestibility, with maximum efficiency at 500 to 1000 phytase units per kg of diet (Jongbloed et al., 2000; Almeida and Stein, 2010). Some studies have also shown an improvement in calcium digestibility as phytase was added to the diet (Adeola et al., 2004; Veum and Ellersieck, 2008; Almeida and Stein, 2010). Improvements in growth rate, feed efficiency, and bone strength (Adeola et al., 2004; Veum and Ellersieck, 2008) and a decrease in phosphorus excretion (Hill et al., 2009) have been reported in pigs fed corn-soybean meal diets supplemented with microbial phytase.

The use of microbial phytase can reduce or eliminate the need for inorganic phosphorus in the diet (Almeida and Stein, 2010) and diet costs are, therefore, often reduced if microbial phytase is used. The digestibility of phosphorus in soybean products is relatively high if microbial phytase is added to the diet; soybean products are, therefore, rich sources of digestible phosphorus, which reduces the need for adding expensive inorganic phosphorus to the diets.

Soybean meal and other soybean products contain relatively high amounts of potassium, magnesium, and sulphur (Table 5), and pigs fed diets containing soybean products and a cereal grain do not need additional supplements of these minerals in their diets. The concentration of micro-minerals may vary according to the geographical area where the soybeans were grown because the concentration of micro-minerals in the soybeans to some degree reflects the concentration of these minerals in the soil, where the beans were grown. It is, therefore, common practice to supplement diets for pigs with the amounts of micro-minerals needed by the animals and disregard the quantities supplied by soybean meal and cereal grains.

Table 5. Concentrations of minerals in soybeans, soybean meal (SBM), and soy protein isolate (as-fed basis)¹

Product	Full-fat soybeans	Dehulled SBM	Nondehulled SBM	Soy protein isolate
Cl, %	0.03	0.49	0.05	0.02
K, %	1.64	2.24	1.96	0.16
Mg, %	0.28	0.27	0.29	0.05
Na, %	0.03	0.08	0.01	0.14
S, %	0.3	0.40	0.39	-
Cu, ppm	16	15.13	17.38	12.9
Fe, ppm	80	98.19	235	15.16
Mn, ppm	30	35.49	40.64	11.9
Se, ppm	0.11	0.27	0.32	0.14
Zn, ppm	39	48.81	50	40.26

¹Values obtained from NRC (2012).

Protein and amino acids

Soybeans are the gold standard of high quality protein for pigs because their amino acid profile complements the amino acid profiles of several cereal grains. In particular, soybean protein is rich in lysine, threonine, and tryptophan (Table 6). These are the most limiting amino acids in corn, wheat, sorghum, and barley (Lewis, 1985). The proteins in soy products are also highly digestible (Table 7).

Table 6. Concentrations of crude protein and amino acids (%) in soybeans, soybean meal (SBM), and other soybean products (as-fed basis)¹

Product	Full-fat soybeans	Dehulled SBM	Nondehulled SBM	Extruded-expelled SBM	Enzyme treated SBM	Fermented SBM	Soy protein concentrate	Soy protein isolate
CP	37.56	47.73	43.90	44.56	55.62	54.07	65.20	84.78
Indispensable AA								
Arg	2.45	3.45	3.17	3.13	3.95	3.70	4.75	6.14
His	0.88	1.28	1.26	1.17	1.41	1.37	1.70	2.19
Ile	1.60	2.14	1.96	1.97	2.48	2.55	2.99	3.83
Leu	2.67	3.62	3.43	3.29	4.09	4.25	5.16	6.76
Lys	2.23	2.96	2.76	2.85	3.20	3.14	4.09	5.19
Met	0.55	0.66	0.60	0.56	0.71	0.75	0.87	1.11
Phe	1.74	2.40	2.26	2.19	2.78	2.87	3.38	4.40
Thr	1.42	1.86	1.76	1.73	2.13	2.09	2.52	3.09
Trp	0.49	0.66	0.59	0.67	0.72	0.69	0.81	1.13
Val	1.73	2.23	1.93	2.06	2.57	2.67	3.14	4.02
Dispensable AA								
Ala	1.59	2.06	1.92	1.89	2.41	2.45	2.82	3.54
Asp	3.89	5.41	4.88	4.84	6.14	5.98	7.58	9.64
Cys	0.59	0.70	0.68	0.70	0.78	0.77	0.90	0.98
Glu	6.05	8.54	7.87	7.56	9.62	9.12	12.02	16.00
Gly	1.52	1.99	1.89	1.89	2.32	2.34	2.75	3.54
Ser	1.67	2.36	2.14	2.11	2.66	2.51	3.33	4.37
Tyr	1.20	1.59	1.55	1.50	2.03	2.08	2.26	3.08

¹Values obtained from NRC (2012).

Table 7. Standardised ileal digestibility (%) of crude protein and amino acids in soybeans, soybean meal (SBM) and other soybean products

Product	Full-fat soybeans	Dehulled SBM	Nondehulled SBM	Extruded-expelled SBM	Enzyme treated SBM	Fermented SBM	Soy protein concentrate	Soy protein isolate
CP	79	87	85	89	88	79	89	89
Indispensable AA								

Arg	87	94	92	96	96	90	95	94
His	81	90	86	91	90	81	91	88
Ile	78	89	88	91	90	81	91	88
Leu	78	88	86	89	89	82	91	89
Lys	81	89	88	90	96	75	91	91
Met	80	90	89	91	91	88	92	86
Phe	79	88	87	90	86	80	90	88
Thr	76	85	83	85	83	73	86	83
Trp	82	91	90	89	83	78	88	87
Val	77	87	84	88	89	80	90	86
Dispensable AA								
Ala	79	85	86	88	86	79	89	90
Asp	80	87	86	88	86	78	88	92
Cys	76	84	84	83	73	64	79	79
Glu	84	89	88	90	88	78	91	94
Gly	81	84	83	84	89	75	88	89
Ser	79	89	89	89	87	80	91	93
Tyr	81	88	86	89	92	88	83	88

¹Values obtained from NRC (2012).

Lysine is the first limiting amino acid in several cereal grains;

therefore, diets are formulated to contain adequate concentrations of lysine. Diets are most correctly formulated based on values for standardized ileal digestibility (SID) of amino acids because values for SID of amino acids are additive in mixed diets (Stein et al., 2007). The SID tryptophan:lysine ratios for soy products range from 19.1 to 23.0 per cent, compared with the ideal ratio of 18.2 per cent (Petersen, 2011).

The concentration of crude protein and amino acids is greater in soybean meal than in full-fat soybeans. Enzyme treatment or fermentation increases the concentration of crude protein in soybean meal because of the removal of oligosaccharides (Cervantes-Pahm and Stein, 2010). The SID for most amino acids in enzyme-treated soybean meal does not differ from that of conventional soybean meal; however, increased concentrations of fat in diets fed to pigs usually increase the digestibility of amino acids because of a reduced rate of passage (Cervantes-Pahm and Stein, 2008).

*
Soybean meal is the premier source of digestible amino acids in diets fed to pigs

Soybean meal is the premier protein source for pigs and is used throughout the world. The reason for the popularity of soybean meal is that the quality of the protein in soybean meal is greater than the quality of protein in other protein meals. Protein quality is often evaluated by expressing the

concentration of each amino acid as a percentage of crude protein. Based on this comparison, it is easily seen that soy protein has greater concentrations of lysine and tryptophan, which are the first limiting amino acids in corn protein when fed to pigs (Table 8). In contrast, corn protein has a greater concentration of methionine than soy protein, which is the reason corn and soybean meal complement each other.

Table 8. Quality of protein in soybean meal and other protein meals expressed as the percentage of each amino acid as a percentage of crude protein (CP)¹

Item	Soybean meal ²		DDGS ³		Canola meal ⁴		Sunflower meal ⁵	
	%	% of CP	%	% of CP	%	% of CP	%	% of CP
Crude protein	47.73	-	27.33	-	37.50	-	39.86	-
Amino acid								
Isoleucine	2.14	4.5	1.02	3.7	1.42	3.8	1.54	3.9
Lysine	2.96	6.2	0.77	2.8	2.07	5.5	1.45	3.6
Methionine	0.66	1.4	0.55	2.0	0.71	1.9	0.78	2.0
Threonine	1.86	3.9	0.99	3.6	1.55	4.1	1.37	3.4
Tryptophan	0.66	1.4	0.21	0.8	0.43	1.1	0.48	1.2
Valine	2.23	4.7	1.35	4.9	1.78	4.7	1.76	4.4

¹Values obtained from NRC (2012).

²Soybean meal, dehulled, solvent extracted.

³DDGS = distillers dried grains with solubles.

⁴Canola meal, solvent extracted.

⁵Sunflower meal, dehulled, solvent extracted.

The requirements for amino acids by pigs are expressed as grams or percentages of digestible amino acids in the diets. These values are calculated by multiplying the concentration of amino acids in feed ingredients by the digestibility of that amino acid. The amino acids provided by soy protein are highly digestible (Figure 4). The concentration of digestible amino acids in soybean meal is greater than in other protein sources that are available to pigs (Figure 5).





Figure 4. Standardised ileal digestibility of amino acids in feed ingredients, % (Values obtained from NRC, 2012)






Figure 5. Digestible amino acids in feed ingredients, g/kg
(Values obtained from NRC, 2012)

One of the challenges in producing diets for pigs is to account for the variability in digestibility that always exists among different batches of the same feed ingredient.

This variability makes it more difficult to predict the concentration of digestible amino acids in the diet that is mixed for the pigs. However, there is less variability in the digestibility of amino acids in soybean meal than in other protein meals. This can be illustrated by calculating the standard deviation of digestibility values in feed ingredients. Such calculations indicate that there is less variability in the digestibility of amino acids among different batches of

soybean meal than among different batches of other protein meals (Table 9). A reduced variability in digestibility values for soybean meal will give users confidence that the desired levels of digestible amino acids in the diets will be achieved if soybean meal is used.

Table 9. Variability in digestibility values for amino acids in soybean meal and other protein meals expressed as standard deviation (SD) for each digestibility value¹

Amino acid	Soybean meal ²		DDGS ³		Canola meal ⁴		Sunflower meal ⁵	
	SID ⁶	SD	SID	SD	SID	SD	SID	SD
Arginine	94	3.12	81	5.25	85	5.56	93	3.35
Histidine	90	4.15	78	4.75	78	10.24	85	6.28
Isoleucine	89	3.79	76	4.87	76	8.34	80	6.15
Leucine	88	3.45	84	4.00	78	6.44	80	5.27
Lysine	89	3.44	61	8.75	74	9.65	78	5.13
Methionine	90	4.70	82	4.13	85	4.06	89	-
Phenylalanine	88	3.65	81	3.96	77	8.42	81	7.11
Threonine	85	4.47	71	5.73	70	9.64	77	8.54
Tryptophan	91	3.32	71	8.16	71	-	80	-
Valine	87	4.16	75	4.95	74	9.78	79	8.06

¹Values obtained from NRC (2012).

²Soybean meal, dehulled, solvent extracted.

³DDGS = distillers dried grains with solubles.

⁴Canola meal, solvent extracted.

⁵Sunflower meal, dehulled, solvent extracted.

⁶SID = standardized ileal digestibility (%).

Fatty acid composition of soybean oil

Soybean oil contains mainly unsaturated fatty acids and less than 15 per cent of all the fatty acids in soybean oil are saturated fatty acids (Table 10). Approximately 50 per cent of all the fatty acids in soybean oil is linoleic acid (C18:2) and an additional 22 per cent is present as monounsaturated fatty acids, primarily oleic acid (C18:1). Soybean oil does, however, also contain more than six per cent linolenic acid (C18:3), which may have anti-inflammatory properties in the diets (NRC, 2012).

Table 10. Fatty acid profile (% of ether extract) and iodine value of full fat soybeans¹

Fatty acid	Abbreviation	%
Myristoleic acid	C-14:0	0.28
Palmitic acid	C-16:0	10.62
Palmitoleic acid	C-16:1	0.28
Stearic acid	C-18:0	3.57
Oleic acid	C-18:1	21.81
Linoleic acid	C-18:2	49.79
Linolenic acid	C-18:3	6.67
Saturated fatty acids	-	14.46
Monounsaturated fatty acids	-	22.09
Polyunsaturated fatty acids	-	56.46
Iodine value	-	128.24

¹Values obtained from NRC (2012).

The high concentration of unsaturated fatty acids in soybean oil may result in deposition of unsaturated fatty acids in back fat and belly fat of pigs fed soybean oil, full fat soybeans or extruded-expelled soybean meal. This may result in increased iodine values in the fat depots, which may create problems in the processing of bellies and loins and potentially reduce the shelf life of these products. It is, therefore, not recommended to feed high levels of soybean oil in diets for pigs during the last few weeks before slaughter.

Usage of Soybean Products

Soybean meal is the most commonly used source of amino acids in diets for pigs around the world. Diets that contain a source of cereal grains, soybean meal and microbial phytase will satisfy the needs for all amino acids, all the energy and most of the phosphorus for growing and finishing pigs and there are usually no needs for other sources of energy and amino acids in these diets.

In diets for weanling pigs, the concentration of conventional soybean meal should be less than 20 per cent to limit the inclusion of antigens and oligosaccharides but fermented soybean meal, enzyme treated soybean meal, or soy protein concentrate can be included in these diets to satisfy the requirement for most of the additional amino acids. Soybean

meal can also satisfy the needs for all amino acids in diets fed to both gestating and lactating sows.

Conclusions

Soybean products are excellent sources of protein for pigs because their amino acid profiles complement those of cereal grains. Amino acids in soy protein are more digestible than amino acids in most other plant proteins, which results in less nitrogen being excreted in the manure from pigs fed diets containing soybean meal than if other protein sources are used.

* Fermented or enzyme treated soybean meal can replace fish meal in diets fed to weanling pigs

Processing of soybean meal to reduce anti-nutritional factors

improves the digestibility of nutrients and energy, and soybean meal that is fermented or enzyme treated may be used in diets fed to weanling pigs as replacement for fishmeal and other animal proteins.

Soybean meal contributes approximately the same quantity of digestible energy to the diets as corn but the energy contribution from full fat soybeans and from extruded-expelled soybean meal is greater than from conventional soybean meal.

Soybean products have a relatively high concentration of phosphorus and if microbial phytase is used in the diet, the digestibility of the phosphorus in soybean products is relatively high. Inclusion of inorganic sources of phosphorus can, therefore, be reduced if soybean products are used together with microbial phytase.

Soybean meal can satisfy the entire requirement for amino acids in diets fed to growing and finishing pigs and to gestating and lactating sows. However, inclusion of conventional soybean meal in diets fed to newly weaned pigs should be restricted to less than 20 per cent but enzyme treated or fermented soybean meal may be used in these diets as additional sources of amino acids.

Literature cited

Adeola, O., J.S. Sands, P.H. Simmins and H. Schulze. 2004. The efficacy of an Escherichia coli-derived phytase preparation. *J. Anim. Sci.* 82:2657-2666.

Almeida, F.N. and H.H. Stein. 2010. Performance and phosphorus balance of pigs fed diets formulated on the basis of values for standardized total tract digestibility of phosphorus. *J. Anim. Sci.* 88:2968-2977.

ASA. American Soybean Association. 2012. Soy Stats: A Reference Guide to Important Soybean Facts and Figures. Am. Soybean Assoc., St. Louis, MO.

Baker, K.M. and H.H. Stein. 2009. Amino acid digestibility and concentration of digestible and metabolizable energy in soybean meal produced from conventional, high-protein, or low-oligosaccharide varieties of soybeans and fed to growing pigs. *J. Anim. Sci.* 87:2282-2290.

Cervantes-Pahm, S.K. and H.H. Stein. 2008. Effect of dietary soybean oil and soybean protein concentrate on the concentration of digestible amino acids in soybean products fed to growing pigs. *J. Anim. Sci.* 86:1841-1849.

Cervantes-Pahm, S.K. and H.H. Stein. 2010. Ileal digestibility of amino acids in conventional, fermented, and enzyme-treated soybean meal and in soy protein isolate, fish meal, and casein fed to weanling pigs. *J. Anim. Sci.* 88:2674-2683.

Cromwell, G.L. 2000. Utilization of soy products in swine diets. Pages 258-282 in *Soy in Animal Nutrition*. J.K. Drackley, ed. Federation of Animal Sciences Societies, Savoy, IL.

Goebel, K.P. and H.H. Stein. 2011. Phosphorus digestibility and energy concentration of enzyme-treated and conventional soybean meal fed to weanling pigs. *J. Anim. Sci.* 89:764-772.

Grieshop, C.M., C.T. Kadzere, G.M. Clapper, E.A. Flickinger, L.L. Bauer, R.L. Frazier and G.C. Fahey. 2003. Chemical and nutritional characteristics of United States soybeans and soybean meals. *J. Agric. Food Chem.*

Hill, B.E., A.L. Sutton and B.T. Richert. 2009. Effects of low-phytic acid corn, low-phytic acid soybean meal, and phytase on nutrient digestibility and excretion in growing pigs. *J. Anim. Sci.* 87:1518-1527.

Jones, C.K., J.M. DeRouchey, J.L. Nelssen, M.D. Tokach, S.S. Dritz and R.D. Goodband. 2010. Effects of fermented soybean meal and specialty animal protein sources on nursery pig performance. *J. Anim. Sci.* 88:1725-1732.

Jongbloed, A.W., P.A. Kemme, Z. Mroz and H.T.M. van Diepen. 2000. Efficacy, use and application of microbial phytase in pig production: A review. Pages 111-129 in *Biotechnology in the Feed Industry*. Alltech's 16th Annu. Symp. T.P. Lyons and K.A. Jacques, ed. Proc. Nottingham Univ. Press, Nottingham, UK.

Kim, S.W., E. van Heugten, F. Ji, C.H. Lee, and R.D. Mateo. 2010. Fermented soybean meal as a vegetable protein source for nursery pigs: I. Effects on growth performance of nursery pigs. *J. Anim. Sci.* 88:214-224.

Lewis, A.J. 1985. Use of synthetic amino acids in practical rations. In *Proceedings of the Carolina Swine Nutrition Conference*, North Carolina State University, Raleigh, 32.

Li, D.F., J.L. Nelssen, P.G. Reddy, F. Blecha, R.D. Klemm, D.W. Giesting, J.D. Hancock, G.L. Allee and R.D. Goodband. 1991. Measuring suitability of soybean products for early-weaned pigs with immunological criteria. *J. Anim. Sci.* 69:3299–3307.

Middelbos, I.S. and G.C. Fahey, Jr. 2008. Soybean carbohydrates. Pages 269-296 in *Soybeans Chemistry, Production Processing, and Utilization*. L.A. Johnson, P.J. White, and R. Galloway, eds. AOCS Press, Urbana, IL.

NRC. National Research Council. 2012. *Nutrient Requirements of Swine*. 11th rev. ed. Natl. Acad. Press, Washington DC.

Petersen, G.I. 2011. Estimation of the ideal standardized ileal digestible tryptophan:lysine ratio in 10 to 20 kg pigs. PhD Diss. Univ. Illinois. Urbana-Champaign.

Rojas, O.J. 2012. Nutritional evaluation of fermented soybean meal fed to weanling pigs. Master's thesis, University of Illinois at Urbana-Champaign.

Sissons, J.W., A. Nyrup, P.J. Kilshaw and R.H. Smith. 1982. Ethanol denaturation of soya bean protein antigens. *J. Sci. Food Agric.* 33:706-710.

Stein, H.H., B. Seve, M.F. Fuller, P.J. Moughan and C.F.M. de Lange. 2007. Invited review: Amino acid bioavailability and digestibility in pig feed ingredients: Terminology and application. *J. Anim. Sci.* 85:172-180.

Wang, T. and L.A. Johnson. 2001. Survey of soybean oil and meal qualities produced by different processes. *J. Am. Oil. Chem. Soc.* 78:311-318.

Veum, T.L. and M.R. Ellersieck. 2008. Effect of low doses of *Aspergillus niger* phytase on growth performance, bone strength, and nutrient absorption and excretion by growing and finishing swine fed corn-soybean meal diets deficient in available phosphorus and calcium. *J. Anim. Sci.* 86:858-870.

July 2013

Jackie Linden

[Learn more](#)
